

# DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

## Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Alcohol Warning: If you consume 3 or more alcohol drinks per day, ask your Doctor whether you should take Ibuprofen or other pain relievers/fever reducers.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if you have

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- taken a diuretic
- reached age 60 or older

Ask a doctor before use if you are

- taking aspirin for heart attack or stroke because ibuprofen may decrease this benefit of aspirin
- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug
- consuming 3 or more alcohol drinks per day

Ask a doctor or pharmacist before use if you are

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use ay increase the risk of heart attack or stroke

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children In case of overdose, get medical help or contact a Poison Control Center 1-800-362-0101 (Arizona only) immediately.

## Directions

- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in a 24-hour period, unless directed by a physician
- take with food or milk, if occasional and mild heartburn, upset stomach, or stomach pain occurs with use
- consult a physician if these symptoms are more thsn mild or if they persist

Children under 12 years

- not for use

## Other information

### Inactive ingredientsOther information

- store at 15 – 30°C (59-86°F).

Inactive ingredients

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Corn Starch, Titanium Dioxide.